

ORIGINAL PAPER

doi: 10.5455/medarh.2016.70.466-469

Med Arch. 2016 Dec; 70(6): 466-469

Received: OCT 25, 2016 | Accepted: DEC 15, 2016

© 2016 Mirjana Gojkov-Vukelic, Sanja Hadzic, Amila Zukanovic, Enes Pasic, Veriva Pavlic

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Application of Diode Laser in the Treatment of Dentine Hypersensitivity

Mirjana Gojkov-Vukelic¹, Sanja Hadzic¹, Amila Zukanovic², Enes Pasic¹, Veriva Pavlic³

¹Department of Oral Medicine and Periodontology, Faculty of Dentistry, University of Sarajevo, Bosnia and Herzegovina

²Department of Pediatric Dentistry, Faculty of Dentistry, University of Sarajevo, Bosnia and Herzegovina

³Department of Periodontology and Oral Medicine, Institute of Dentistry, Banja Luka, Bosnia and Herzegovina

Corresponding author: Mirjana Gojkov Vukelic, DMD, PhD. Department of Oral Medicine and Periodontology, Faculty of Dentistry, Sarajevo. Bolnička 4a, 71000 Sarajevo, Bosnia and Herzegovina. Tel: +387 33 407 863. E-mail address: gojkov@bih.net.ba

ABSTRACT

Introduction: Dentine hypersensitivity is characterized by acute, sharp pain arising from the exposed dentine, most commonly in response to thermal, tactile, or chemical stimuli, and which cannot be linked to any other pathological changes in the tooth or the environment. Therapy uses various impregnating agents in the form of solutions or gels and, in more recent times, laser. **Aim:** The aim of this research was to examine the effects of treatment of hypersensitive dental cervix with diode laser. **Materials and Methods:** The study included 18 patients with 82 sensitive teeth. The degree of dentine hypersensitivity was evaluated by visual analogue scale (VAS), and the treatment was carried out by application of low-power diode laser over the span of three visits, which depended on the initial sensitivity. **Results:** There is a significant difference in VAS values measured at the onset of treatment (baseline) and immediately after the first laser treatment ($t=9.275$; $p=0.000$), after 7 days, after the second laser treatment (14 days) ($t=7.085$, $p=0.000$), as well as after 14 days and the third laser treatment ($t=5.517$, $p=0.000$), which confirms the effectiveness of this therapeutic procedure. The results showed a reduction of hypersensitivity in response to tactile stimulus with a probe after the third treatment, even with teeth whose value on the VAS was very high at the beginning of treatment (baseline). **Conclusion:** Within the scope of the conducted study, laser therapy has provided extremely safe and effective results in the treatment of cervical dentine hypersensitivity.

Keywords: dentine hypersensitivity, desensitizing agent, diode laser, laser therapy.

1. INTRODUCTION

Thanks to good preventive dental programs and developed knowledge about the importance of oral hygiene, the vitality of teeth within the oral cavity has been extended in the recent times, which led to an increase in non-carious cervical lesions, or dental erosions, abrasions, etc. (1, 2).

Dentine hypersensitivity (DH) is characterized by acute, sharp pain in the area of exposed dentine, in response to thermal, chemical, osmotic, and tactile stimuli (3). Although sensitivity can occur on any part of the tooth, it is most commonly felt in the vestibular area of dental cervical region (for canines and first premo-

lars) and on the root surface. The frequency at which it occurs ranges between 3 and 57%, and it is much more frequent (72-98%) in patients suffering from periodontal disease. It most often occurs between 20 and 50 years of age, and is more common among women (4, 5).

Difficulties in treating cervical DH gave rise to a large number of techniques and therapeutic procedures which are currently used for pain alleviation in DH (6).

Therapy uses various impregnating agents in the form of solutions or gels and, in more recent times, laser. Based on hydrodynamic theory, several methods, such as the application of fluoride, dentine adhesives, corti-

Number of treatments	Number of teeth N	Percentage %	X VAS base-line	SD
1	27	32.92	2.90	±2.09
2	29	35.36	4.36	±2.49
3	26	31.70	4.59	±2.73
Total	82	100	3.95	±2.53

Table 1. Number of treatments compared to average VAS value at baseline (at the beginning of treatment)

	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	45.54	2	22.77	3.77	0.027
Within groups	476.13	79	6.02		
Total	521.68	81			

Table 2. Baseline assessment of tooth sensitivity and number of treatments

costeroids, and silver nitrate solution, work by blocking the open dentine tubules.

In recent decades, classic treatments with desensitizing agents have been supplemented by the use of laser. Using lasers to treat DH dates back to the 80s with the advent of erbium lasers. Even though the initial results were quite disappointing, the improvement of technology and scientific knowledge over time has led to the development of new lasers with wavelengths suitable for therapeutic treatment (7).

Recent studies report satisfactory results of treatment with laser irradiation. Most studies conducted with different types of lasers, with different wavelengths and duration of application, reveal the effectiveness of this treatment, both immediately upon the completion of therapy, and after circa 6 months from the first treatment. As a result, the pain is reduced and, in many cases, eradicated (8, 9, 10). Among the published works, there are those which confirm the exceptional efficacy of the use of diode lasers in the treatment of DH. Thus, the aim of our study was to investigate the effects of diode laser therapy on hypersensitive dental cervix.

2. MATERIALS AND METHODS

The study included 18 patients with 82 sensitive teeth. The degree of dentine hypersensitivity was evaluated by visual analogue scale (VAS). Dentine hypersensitivity was stimulated by touching the dental cervix with the tip of the probe, with mesial-distal directionality. All patients were asked to assess their level of dentine hypersensitivity using the VAS scale of 0 to 10, where 0 represents “no pain” and 10 represents “greatest pain.”

After initial sensitivity was assessed and recorded, laser therapy was initiated.

Laser treatment protocol: Low-power diode laser (SmilePro980, Biolitec) was used in this study. The laser was operated in a continual regime, and 2 W of power was applied to the tooth surface. During the 60 seconds of exposure, tooth tissue was around 2mm away from the laser.

Exposure time (60s) was repeated after sensitivity control (using the VAS scale), seven and fourteen days after initial exposure, only on those teeth that were still sensitive. While working with the laser, both the therapist and the patient wore protective goggles, and work space was appropriately designated and marked.

3. RESULTS

The study included 18 patients, with average age of 27 years, who had 82 sensitive teeth.

It can be seen that initially less sensitive teeth required fewer treatments!

ANOVA test was carried out in order to assess whether this difference is significant.

There is a significant difference in tooth sensitivity values measured at baseline, in teeth that had a different number of laser treatments. ANOVA $F=3.77$, $p=0.027$.

Based on the obtained results, we can say (with 95% confidence) that teeth which had lower dentine sensitivity at the very beginning will require fewer laser treatments.

In order to determine between which teeth this difference is observed, given the number of treatments, a post-hoc analysis was carried out using Turkey’s Honest Significant Difference (HSD) test.

Differences occur only between the mean sensitivity values at baseline between teeth that had only one treatment and teeth that had three laser treatments ($p=0.037$), but there is no difference between VAS value at baseline between teeth that had one laser treatment and those that were treated twice ($p=0.073$), nor between the teeth that had two and those that had three laser treatments ($p=0.934$).

There is a significant difference in VAS values measured at baseline and after the first laser treatment: $t=9.275$, $p=0.000$. There is a significant difference in VAS values measured at baseline and after the second laser treatment: $t= 1.268$, $p=0.000$. There is a significant difference in VAS values measured at baseline and after the third laser treatment: $t=8.749$, $p=0.000$.

	Paired differences					t	df	Sig. (2-tailed)
	Mean	Std. Deviation	Std. error mean	95% Confidence Interval of the difference				
				Lower	Upper			
Baseline tooth sensitivity assessment – Immediate tooth sensitivity following the first treatment	1.97	1.92	0.21	1.55	2.39	9.27	81	0.000
Baseline tooth sensitivity assessment – Tooth sensitivity following the second treatment (7 days)	3.19	2.53	0.28	2.62	3.75	11.26	79	0.000
Baseline tooth sensitivity assessment – Tooth sensitivity following the second treatment (14 days)	3.70	2.71	0.42	2.85	4.56	8.74	40	0.000

Table 3. VAS values of all teeth before treatment and following the first, second, and third laser application

	Paired differences					t	df	Sig. (2-tailed)
	Mean	Std. Devi- ation	Std. error mean	95% Confidence In- terval of the difference				
				Lower	Upper			
Baseline tooth sensitivity assessment - Immediate tooth sensitivity following the first treatment	1.97	1.92	0.21	1.55	2.39	9.27	81	0.000
Immediate tooth sensitivity following the first treatment - Baseline assess- ment (7 days)	0.09	2.12	0.23	-0.38	0.56	0.37	79	0.706
Baseline assessment - Tooth sensi- tivity following the second treatment (7 days)	1.13	1.43	0.16	0.81	1.45	7.08	79	0.000
Tooth sensitivity following the second treatment - Baseline assessment (14 days)	-0.08	1.59	0.24	-0.58	0.41	-0.34	40	0.733
Baseline assessment - Tooth sensi- tivity following the second treatment (14 days)	0.74	0.86	0.13	0.46	1.01	5.51	40	0.000
Baseline tooth sensitivity assessment - Tooth sensitivity following the second treatment (7 days)	3.19	2.53	0.28	2.62	3.75	11.26	79	0.000
Baseline tooth sensitivity assessment - Tooth sensitivity following the second treatment (14 days)	3.70	2.71	0.42	2.85	4.56	8.74	40	0.000

Table 4. VAS values for all teeth after the first, second, and third application of laser, and baseline measurement prior to the application of laser

There is a significant difference in VAS values measured at baseline and after the first laser treatment ($t=9.275$; $p=0.000$), as well as after 7 days and after the second laser treatment (14 days) ($t=7.085$, $p=0.000$), and after 14 days and the third laser treatment ($t=.517$, $p=0.000$), which supports the effectiveness of this therapeutic procedure.

4. DISCUSSION

Dentine hypersensitivity (DH) is common, and individual needs for treatment depend on aetiology, as well as on the subjective experience of painful sensations and the degree of tolerance to this type of pain.

In this study, some of the patients reported pain so severe that it has become a physical and emotional problem that affects their quality of life. Many of them were not able to consume hot or cold foods or liquids, acidic foods or liquids, and even had difficulty with brushing teeth. As the data from previous studies suggest, several methods should be applied during treatment in order to obtain satisfactory results, since the aetiology of DH may be multifactorial (11, 12, 13).

Conventional methods of treating DH include topical application of desensitizing agents, either professionally or at home, such as protein precipitates, agents for occlusion of dentinal tubules (14) and, more recently, lasers (15, 16, 17). It is believed that the occlusion of dentinal tubules leads to a decrease in permeability of dentine and, proportionally, also reduces DH (18). According to hydrodynamic theory, efficacy of dentine desensitization agents is directly related to their ability to efficiently close dentinal tubules (19, 20).

In their study, Yilmaz et al. compared the effectiveness of application of sodium fluoride and diode laser in the treatment of DH. They concluded that, within the scope of their study, GaAlAs laser therapy is effective in the

treatment of DH, and is a more comfortable and faster treatment than traditional treatments for DH (21). These results support the results of our study.

Several studies (22, 23) describe the synergistic effect of lasers in conjunction with desensitization agents. For this reason, our study included laser irradiation of the cervical portion of the tooth only, and we obtained exceptionally good results in terms of lowered dentine hypersensitivity ($F = 3.77$, $p = 0.027$). Therefore, we can state (with 95% confidence) that teeth which had lower dentine sensitivity at the very beginning will require fewer laser treatments.

Previous published data indicate that only the Nd:YAG laser has an additional analgesic effect, compared with other lasers. These findings are the result of the effect of radiation which can temporarily alter the endings of sensory axons and block both C and AB fibres, thereby reducing the pain (24).

Parameter of the power used in our study was 2 W, which is in accordance with the study by Liu et al. (25). Their study (25) demonstrated that 2 W (166 J/cm^2) is a suitable parameter for the 980nm diode laser, which sealed dentinal tubules without excessive melting of the dentine, thus achieving a good level of analgesia, which is comparable to our results. Good results arise from the closure of dentinal tubules, which prevents internal communication of dental pulp with oral cavity fluids (15, 26).

Based on the results of our study in which only a diode laser was used, we believe that modern low-power lasers can also provide good results in the treatment of DH; this finding is also supported by the results of research by Umberto et al. (27).

Our research, as well as research by other authors (28, 29), demonstrates that low-energy lasers, including GaAlAs diode laser with wavelengths between 780 and

980 nm, have an effect on nerve endings, thus eliminating sensitivity.

In a study conducted on 27 patients with 55 hypersensitive teeth, Lopes et al. assessed the efficacy of various protocols for treating dentine hypersensitivity. They concluded that all desensitising protocols are effective in reducing dentine hypersensitivity, but have different effects. Therefore, they believe that a combination of protocols is an interesting alternative for the treatment of cervical dentine hypersensitivity (30). This conclusion follows from the need to achieve satisfactory results in as few treatments as possible. The results of our study indicate that, applied through multiple treatments, this modern therapeutic procedure independently achieves good results, even in teeth with greater level of hypersensitivity.

We believe that further research is needed to assess long-term effects of these therapeutic procedures on a larger sample in order to provide recommendations for use in routine clinical practice.

5. CONCLUSION

Within the scope of the conducted study, laser therapy has provided extremely safe and effective results in the treatment of cervical dentine hypersensitivity.

- Authors' contributions: Conception and design: MGV, EP and VP; Acquisition, analysis and interpretation of data: SH, EP and AZ; Drafting the article: MGV and AZ; Revising it critically for important intellectual content: MGV, VP, SH.

REFERENCES

1. Bartlett DW. The role of erosion in tooth wear: aetiology, prevention and management. *Int Dent J.* 2005; 55: 277-84.
2. Bartlett D.W. Shah P. A critical review of non-carious cervical (wear) lesions and the role of abfraction, erosion, and abrasion. *J Dent Res.* 2006; 85: 306-12.
3. Ady M. Dentine hypersensitivity. New perspectives and old problem. *Inter Dent J.* 2002; 52: 367- 78.
4. Rees JS. The prevalence of dentine hypersensitivity in general dental practise in the UK. *J Clin Periodontol.* 2000; 27: 860-5.
5. Walline BW, Wagner JG, Marx DB, Reinhart RA. Comparison of methods for measuring root and mucogingival sensitivity. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2000; 90(5): 641-6.
6. Porto ICCM, Andrade AKM, Montes MAJR. Diagnosis and treatment of dentinal hypersensitivity. *J Oral Sci.* 2009; 51: 323-32.
7. Bamise CT, Esan TA. Mechanisms and treatment approaches of dentine hypersensitivity: a literature review. *Oral Health Prev Dent.* 2011; 9: 353-67.
8. Blatz MB. Laser therapy may be better than topical desensitizing agents for treating dental hypersensitivity. *J Evid Based Dent Pract.* 2012; 12: 69-70.
9. da Rosa WL, Lund RG, Piva E, da Silva AF. The effectiveness of current dentin desensitizing agents used to treat dental hypersensitivity: a systematic review. *Quintessence Int.* 2013; 44: 535-46.
10. Orchardson R. Gillam DG. Managing dentin hypersensitivity. *J Am Dent Assoc.* 2006; 137: 990-8.
11. Birang R, Poursamimi J, Gutknecht N, Lampert F, Mir M. Comparative evaluation of the effects of Nd: YAG and Er:YAG laser in dentin hypersensitivity treatment. *Lasers Med Sci.* 2007; 22: 21-4.
12. Kara C, Orbak R. Comparative evaluation of Nd: YAG laser and fluoride varnish for the treatment of dentinal hypersensitivity. *J Endod.* 2009; 35: 971-4.
13. Madhavan S, Nayak M, Shenoy A, Shetty R, Prasad K. Dentinal hypersensitivity: A comparative clinical evaluation of CPP-ACP F, sodium fluoride, propolis, and placebo. *J Conserv Dent.* 2012; 15: 315-8.
14. Gangarosa LP, Sr Current strategies for dentist-applied treatment in the management of hypersensitive dentine. *Arch Oral Biol.* 1994; 39(Suppl): 101S-6S.
15. Kimura Y, Wilder-Smith P, Yonaga K, Matsumoto K. Treatment of dentine hypersensitivity by lasers: A review. *J Clin Periodontol.* 2000; 27: 715-21.
16. He S, Wang Y, Li X, Hu D. Effectiveness of laser therapy and topical desensitizing agents in treating dentine hypersensitivity: a systematic review. *J Oral Rehabil.* 2011; 38: 348-58.
17. Biagi R, Cossellu G, Sarcina M, Pizzamiglio IT, Farronato G. Laser-assisted treatment of dentinal hypersensitivity: a literature review. *Ann Stomatol (Roma).* 2016.
18. Pashley DH. Dentin permeability and dentin sensitivity. *Proc Finn Dent Soc.* 1992; 88 (Suppl 1): 31-7.
19. Bartold PM. Dentinal hypersensitivity: A review. *Aust Dent J.* 2006; 51: 212-8.
20. Walters PA. Dentinal hypersensitivity: A review. *J Contemp Dent Pract.* 2005; 6: 1-10.
21. Yilmaz HG, Kurtulmus-Yilmaz S, Cengiz E. Long-term effect of diode laser irradiation compared to sodium fluoride varnish in the treatment of dentine hypersensitivity in periodontal maintenance patients: a randomized controlled clinical study. *Photomed Laser Surg.* 2011 Nov; 29(11): 721-5.
22. Dilsiz A, Aydin T, Canakci V, Gungormus M. Clinical evaluation of Er: YAG, Nd: YAG, and diode laser therapy for desensitization of teeth with gingival recession. *Photomed Laser Surg.* 2010; 28: S1-S7.
23. Dantas EM, Amorim FK, Nóbrega FJ, Dantas PM, Vasconcelos RG, Queiroz LM, Clinical Efficacy of Fluoride Varnish and Low-Level Laser Radiation in Treating Dentin Hypersensitivity. *Braz Dent J.* 2016 Jan-Feb; 27(1): 79-82.
24. Myers TD, McDaniel JD. The pulsed Nd: YAG dental laser: review of clinical applications. *J Calif Dent Assoc.* 1991; 19: 25-30.
25. Liu Y, Gao J, Gao Y, Xu S, Zhan X, Wu B. In vitro study of dentin hypersensitivity treated by 980-nm diode laser. *J Lasers Med Sci.* 2013; 4: 111-9.
26. Lan WH, Lee BS, Liu HC, Lin CP. Morphologic study of Nd: YAG laser usage in treatment of dentinal hypersensitivity. *J Endod.* 2004; 30: 131-4.
27. Umberto R, Claudia R, Gaspare P, Gianluca T, Alessandro del V. Treatment of dentine hypersensitivity by diode laser: a clinical study. *Int J Dent.* 2012; 2012: 858950.
28. Asnaashari M, Moeini M. Effectiveness of lasers in the treatment of dentin hypersensitivity. *J Lasers Med Sci.* 2013; 4: 1-7.
29. Suri I, Singh P, Shakir QJ, Shetty A, Bapat R, Thakur R. A comparative evaluation to assess the efficacy of 5% sodium fluoride varnish and diode laser and their combined application in the treatment of dentin hypersensitivity. *J Indian Soc Periodontol.* 2016 May-Jun; 20(3): 307-14.
30. Lopes AO, Eduardo Cde P, Aranha AC. Clinical evaluation of low-power laser and a desensitizing agent on dentin hypersensitivity. *Lasers Med Sci.* 2015 Feb; 30(2): 823-9.