

Mat Soc Med. 2012 Dec 24(4): 274-276

Received: 27 September 2012

Accepted: 19 November 2012

Conflict of interest: none declared.

© AVICENA 2012

DOI: 10.5455/msm.2012.24.274-276

Smoking and Periodontal Disease

Tea Borojevic

Pliva, Zagreb, Croatia

Corresponding author: Tea Borojevic, MDD, Pliva, Zagreb, Croatia. E-mail: tea5bor@yahoo.com

REVIEW

ABSTRACT

Periodontitis is a group of inflammatory diseases affecting the supporting tissues of the tooth (periodontium). The periodontium consists of four tissues : gingiva, alveolar bone and periodontal ligaments. Tobacco use is one of the modifiable risk factors and has enormous influence on the development, progres and tretmen results of periodontal disease. The relationship between smoking and periodontal health was investigated as early as the miiddle of last century. Smoking is an independent risk factor for the initiation, extent and severity of periodontal disease. Additionally, smoking can lower the chances for successful tretment. Tretmans in patients with periodontal disease must be focused on understanding the relationship between genetic and environmental factors. Only with individual approach we can identify our pacients risks and achieve better results.

Key words: smoking, periodontal disease.

1. INTRODUCTION

Periodontitis is a group of inflammatory diseases affecting the supporting tissues of the tooth (periodontium). The periodontium consists of four tissues : gingiva, alveolar bone and periodontal ligaments.

The periodontal diseases are highly prevalent and can affect up to 90% of the world wide population. Gingivitis, the mildest form of periodontal disease, is caused by the bacterial biofilm (dental plaque) that accumulates on teeth adjacent to the gingiva (gums).

The simtoms are usualy red, swollen gums who can bleed easily. However, gingivitis does not affect the underlying supporting structures of the teeth and is reversible. When gingivitis is not treated, it can advance to periodontitis. Periodontitis results in loss of connective tissue and bone support and is a major cause of tooth loss in adults (1). In addition to pathogenic microorganisms in the biofilm, genetic and environmental factors has enormouse influence on development periodontal disease. Tobacco use is one of the modifiable risk factors and has enormous influence on the development ,progres and tretmen results of periodontal disease.

The American Academy of Periodontology (AAP) has classified periodontitis into aggressive periodontitis (AgP), chronic periodontitis (CP) and periodontitis as a manifestation of systemic diseases (2). Both AgP and CP have a multi-factorial etiology with dental plaque as the initiating factor (3). However the initiation and progression of periodontitis are influenced by other factors including tobacco use.

2. RELATIONSHIP BETWEEN SMOKING AND PERIODONTAL DISEASE

One-third of the world's adult population are smokers (57% of these are men, 43% are women) . It is predicted that in 20 years this yearly death rate from tobacco use will be more than 10 million people. Smoking in developing countries is rising by more than 3% a year (4). We can assume periodontal diseasees will also rise.

The relationship between smoking and periodontal health was investigated as early as the miiddle of last century. Smoking is an independent risk factor for the initiation, extent and severity of periodontal disease. Additionally, smoking can lower the chances for successful tretment.



Figure 1. Generalized advanced chronic periodontitis in smooker



Figure 2. Generalized advanced chronic periodontitis in smoker

Cross-sectional and longitudinal data provide strong support for the statement that the risk of developing periodontal disease as measured by clinical attachment loss and alveolar bone loss increases with increased smoking. Studies find that former smokers (clinically defined as two or more years since quitting smoking) experience less attachment loss than current smokers but more than never-smokers. Furthermore, the likelihood of developing increasing periodontal disease exhibits dose dependency (5)

For many years sciences did not know how smoking

sight into the established clinical conundrum of increased infection with periodontal pathogens but reduced clinical inflammation in smokers.

Bagaikar and asos. established that exposure of *P. gingivalis* to tobacco smoke extract increased the expression of the major fimbrial antigen (FimA), but not the minor fimbrial antigen (Mfa1). That means that exposure did not induce *P. gingivalis* auto-aggregation but did promote dual species biofilm formation, monitored by microcolony numbers and depth. Interestingly, *P. gingivalis* biofilms grown in the presence of tobacco smoke exhibited a lower pro-inflammatory capacity (TNF- α , IL-6) than control biofilms. The underlying mechanisms are unknown, more likely tobacco smoke represents an environmental stress to which *P. gingivalis* adapts by altering the expression of several virulence factors—including major and minor fimbrial antigens (FimA and Mfa1, respectively) and capsule—concomitant with a reduced pro-inflammatory potential of intact *P. gingivalis*(7,8).

In vitro studies have shown altered gingival crevicular fluid inflammatory cytokine profiles (GCF), immune cell function and altered proteolytic regulation in smokers.

Smokers exhibited a decrease in several pro-inflammatory cytokines and chemokines and certain regulators of T-cells and NK-cells. This reflects the immunosup-




Figure 3. Progression of periodontal disease :<http://www.stcatherinesdentalpractice.co.uk/dental-treatments-in-grantham/periodontal-treatment-in-grantham.html>

affect periodont and why people with chronic periodontitis have reduced clinical inflammation. Today we know that tobacco smoke induces alterations to the 3-OH fatty acids present in lipid A in a manner consistent with a microflora of reduced inflammatory potential.

In investigation smokers had significant reductions in the 3-OH fatty acids associated with the consensus (high potency) enteric LPS structure were noted in smokers compared with non-smokers with chronic periodontitis. Thus, smoking is associated with specific structural alterations to the lipid-A-derived 3-OH fatty acid profile in saliva that are consistent with an oral microflora of reduced inflammatory potential.(6)

These findings provide much-needed mechanistic in-

pressant effects of smoking which may contribute to an enhanced susceptibility to periodontitis (9).

Periodontal treatment tends to be less likely to be successful in smokers than in non-smokers. Studies evaluating the efficacy of periodontal disease control and specific periodontal procedures including regenerative procedures, soft tissue grafting procedures and implant procedures have consistently demonstrated a negative effect from smoking on success rates(10).

3. CONCLUSION

Smoking is well-established risk factor for periodontal disease. It changes the human microflora, human immune response that leads to destruction of the support-

ing tissues of the tooth. Difficult circumstance is a fact that symptoms in periodontal disease in smokers are increased so It can take years before the patient seeks help, then it is often too late.

Tretmans in patients with periodontal disease must be focused on understanding the relationship between genetic and environmental factors. Only with individual approach we can identify our patients risks and achieve better results.

REFERENCES

1. Armitage GC. Development of a classification system for periodontal disease and conditions. *Ann Periodontol*. 1999; 4: 1-6. doi: 10.1902/annals.1999.4.1.1. Page RC, Offenbacher S, Schroeder HE, Seymour GJ, Kornman KS. Advances in the pathogenesis of periodontitis: summary of developments, clinical implications and future directions. *Periodontol* 2000. 1997; 14: 216-248. doi: 10.1111/j.1600-0757.1997.tb00199.x.
2. Pihlstrom BL, Michalowicz BS, Johnson NW. Periodontal disease. *The Lancet*. 2005; 366(9499): 1809-1820.
3. Bagaitkar J, Daep CA, Patel CK, Renaud DE, Demuth DR, Scott DA. Tobacco smoke augments Porphyromonas gingivalis–Streptococcus gordonii biofilm formation. *PLoS One*. 2011; 6(11): e27386.
4. Hayman L, Steffen MJ, Stevens J. Smoking and periodontal disease: discrimination of antibody responses to pathogenic and commensal oral bacteria. *Clin Exp Immunol*. 2011; 164(1): 118-126.
5. Buduneli N, Larsson L, Biyikoglu B, Renaud DE, Bagaitkar J, Scott DA. Fatty acid profiles in smokers with chronic periodontitis. *J Dent Res*. 2011; 90(1): 47-52.
6. <http://www.medicalnewstoday.com/releases/9702.php>
7. <http://dentalstudynotes.pgpreparation.in/2007/12/periodontitis.html>
8. Tomar SL, Asma S. Smoking-attributable periodontitis in the United States: findings from NHANES III. *National Health and Nutrition Examination Survey*. *J Periodontol*. 2000; 71(5): 743-751.
9. Tymkiw KD, Thunell DH, Johnson GK, Joly S, Burnell KK, Cavanaugh JE, Brogden KA, Guthmiller JM. Influence of smoking on gingival crevicular fluid cytokines in severe chronic periodontitis *J Clin Periodontol*. 2011 Mar; 38(3): 219-228. doi: 10.1111/j.1600-051X.2010.01684.x.
10. Bagaitkar J, Demuth DR, Daep CA, Renaud DE, Pierce DL, Scott DA P. gingivalis fimbrial proteins which induce TLR2 hyposensitivity. *PLoS One*. 2010; 5(5): e9323..

Home Login Contact



An independent scientific and professional voice for public health in Europe

- About EUPHA
- Public Health in Europe
- Members
- EUPHA Sections
- EUPHA Conferences
- European Journal of Public Health
- EUPHA Publications
- EUPHApedia
- Projects
- Links

Login to EUPHA

Login:

Password:

Submit

[Forgot Password?](#)
[New User?](#)

The European Public Health Association

The European Public Health Association, or EUPHA in short, is an umbrella organisation for public health associations in Europe. EUPHA was founded in 1992. EUPHA is an international, multidisciplinary, scientific organisation, bringing together around 14000 public health professionals for professional exchange and collaboration throughout Europe. We encourage a multidisciplinary approach to public health.

[Read More](#)

Conferences



EUROPEAN PUBLIC HEALTH CONFERENCE

The next EUPHA conference for 2012 is themed 'All Inclusive Public Health' and will be held in St. Julian's, Malta

In partnership with:

- Association of Schools of Public Health in the European Region (ASPHER)
- Malta Association of Public Health Medicine (MAPHM)

[Read More](#)

News

14 September, 2012
WHO/EURO Regional Committee adopted the **European Action Plan** for Strengthening Public Health Capacities and Services at its 62nd session that concluded yesterday in Malta. [Read more...](#)

05 September, 2012
The EUPHA August newsletter is distributed to the newsletter subscribers. If you are interested, [login to the database](#) and set your newsletter box to YES. [Read more...](#)

03 September, 2012
EUPHA to attend the annual meeting of the German Society of Social Medicine and Prevention! [Read more...](#)

05 July, 2012
Going down memory lane - 20 years of EUPHA in pictures
[Click here](#) for the latest addition. [Read more...](#)

29 June, 2012

276

Review • Mat Soc Med. 2012 Dec 24(4): 274-276